



a division of Amgrow

in focus | THIRAM DG

in focus

THIRAM DG FUNGICIDE

Include Barmac's Thiram DG Fungicide in your strawberry fungicide program this season

**New strawberry registration with 2 day withholding period!
Allows its use during the picking season!**



Thiram DG offers disease protection from:

- **Grey Mould**
- **Black Spot**

Efficacy on Grey Mould (*B. cinerea*) and Black Spot (*Colletotrichum* sp.)

Assessments (days after last application)	Grey Mould			Black Spot	
	1st (0-7)	2nd (8-14)	3rd (15-21)	1st (4-7)	2nd (7-37)
Thiram DG (% effectiveness)	72.26	74.58	74.48	92.1	73.4
Euparen (% effectiveness)	77.62	73.99	67.88	94.7	77.4

Results based on an average of 20 trials.
High level efficacy with Thiram DG, comparable to other specific active ingredients.

Don't let Grey Mould or Black Spot disease bring your strawberry's down!

Situation	Pest	Rate	Comments
Strawberries (QLD, VIC, SA, TAS, WA only).	Black Spot (<i>Colletotrichum actatum</i>), Grey Mould (<i>Botrytis cinerea</i>)	150g/100 litres of water	Apply as a protective spray at flowering then at 10-14 day intervals as necessary until 2 days before harvest. Do Not exceed a maximum water volume of 1200L/Ha.