

ANALYSIS (% w/v):

11.5% Total Nitrogen

4.6% (N) as Urea

6.87% (N) as Ammonium

10.5% Phosphate

4% Potassium

0.35% Zinc

0.13% Boron

2.5% Sugars

7.5% Organic acid

225 mg/L Vitamin B1

15 mg/L Auxins

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 **NutraFeed**[®]
Liquid Fertiliser

Establish

PRODUCT

CHARACTERISTICS:

Colour: Brown

SG: 1.3



a division of Amgrow

Barmac ABN 21 009 674 953

A division of Amgrow

82 Christensen Road,

Stapylton QLD 4207

TEL: (07) 3802 5050

www.barmac.com.au

■ 20L ■ 200L ■ 1000L

Establish is a new effective starter fertiliser suitable for soil application on a wide range of horticultural and broad acre crops.



Directions for use

Suitable for use via fertigation

Crop	Fertigation (L/ha)	Comments
Berry fruits	8-10	Apply weekly after planting for the first month
Citrus	10-15	Apply fortnightly after planting for the first three months
Cucurbits	8-10	Apply weekly after planting for the first month
Broadacre (cotton beans, corn)	5-7	Apply via water injection or directed furrow spray at planting or after emergence
Nursery/ Ornamental	1:100	Drench trays/pots/tables with a 1:100 dilution solution
Pomefruit / Stonefruit	10-15	Apply fortnightly after planting for the first three months
Potato	8-10	Apply weekly via overhead irrigation for the first month or as a directed furrow spray at planting
Pasture Establishment	10	Foliar apply after sowing and irrigate into soil
Sugarcane	4-5	Apply to billets at planting via spray or dip
Tomato/Capsicum	8-10	Apply weekly after planting for the first month
Tree Nut Orchards	10-15	Apply fortnightly after planting for the first three months. Apply after tree harvest shaking to assist root anchoring
Tropical Fruit (avocado, mango, banana, pines etc)	10-15	Apply fortnightly after planting for the first three months
Turf	20	Apply weekly after sowing/renovating for the first month

Vegetables / Brassica	8-10	Apply weekly after planting for the first month
Vines / Grapes	8-10	Apply fortnightly after planting for the first three months

CONDITIONS OF SALE - The suggested rates of application are designed for typical use conditions and should be used as a guide only. Because climatic and soil conditions, application methods, irrigation and agricultural practices are beyond the control of Barmac and cannot be foreseen, Barmac accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.

STORAGE AND HANDLING: Store in a cool, well-ventilated area, out of direct sunlight, avoiding temperature extremes. Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry, locked area out of reach of children. Protect from excessive heat or cold. Triple rinse (or equivalent) and dispose of container in accordance with local, state and federal regulations.

WARNING: Avoid skin and eye contact, ingestion and contact with or inhalation of spraymist. When handling this product, wear suitable gloves, goggles, hat and protective clothing.

FIRST AID: In case of contact with eyes or skin, flush at once with large amounts of water. If irritation persists, seek medical attention. If swallowed, drink large amounts of water and induce vomiting. Seek medical attention.

BATCH:

DATE OF MANUFACTURE: