

## Double Time

Active Constituents: 340g/L MCPA and 80g/L Dicamba

For the post emergent control of many broadleaf weeds in winter cereals, grass pastures and turf situations excluding buffalo grass.

### Double Time Benefits:

- A safe broad spectrum broadleaf herbicide that is suitable for use on cereals, grass pastures and turf to bring weed pressure under control
- Economically and safely controls many broadleaf weeds in registered crops
- Quick control with weeds knocked over within 7-14 days
- Easy to use and apply and can be tank mixed with compatible pesticides and fertilisers



Capeweed in Wheat



Wild Radish in Cereal Crop

### Product Characteristics:

Colour	Specific Gravity
Brown Liquid	1.11

**Pack Sizes:** Available in 1L, 5L and 20L packs



a division of  
Amgrow Australia Pty Ltd

# Double Time - Post Emergent Selective Herbicide

**Application: (Refer to product label for more detailed instructions)**

CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
<b>Barley, Commercial Rye, Oats, Triticale, Wheat</b>  From early tillering stage (Zadoks 21)	Capeweed, Charlock, Common Ice-Plant, seedling Docks, Doublegee (Three cornered Jack, Spiny Emex), Hexham Scent (Melilotus), Mustards, Soldier Thistle, Sorrel Seedlings, Tares, Tree Hogweed, Turnip Weed (Rapistrum rugosum), Vetches, Ward's Weed, Wild Radish, Wild Turnip (Brassica tournefortii), Wireweed (Hogweed)	All States	1 Litre per hectare	Spry when most weeds are in the 2 to 4 leaf stage. In some instances suppression only of Wireweed (Hogweed), Capeweed and Wild Radish will occur
	Seedling Cotton Thistle, Fat Hen	TAS Only		
<b>Barley, Commercial Rye, Oats, Triticale, Wheat</b>  From early tillering stage (Zadoks 21) until the fully tillered stage and before joining occurs (Zadoks 30)	Cotton thistle to bud formation	TAS Only	1.7 Litres per hectare	Spray when most weeds are still in the seedling stage. Note: Wild Radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances, suppression only of Wireweed (Hogweed), Hoary Cress and Climbing Buckwheat will be obtained.
	Mintweed, New Zealand Spinach	QLD, NSW ACT Only		
	Capeweed, Charlock, Chickweed, Climbing Buckwheat (Black Bindweed), Clover, Common Ice-Plant, seedling Docks, Doublegee (Three-cornered Jack), Spiny Emex), Fat-Hen, Hexham Scent (Melilotus), seedling Hoary Cress, Mouse-eared Chickweed, Mustards, Narrow-leaved Plantain, Saffron Thistle, Spurry, Swinecress, Tares, Tree Hogweed, Turnip Weed (Rapistrum rugosum), Variegated Thistle, Vetch, Wild Radish, Wild Turnip (Brassica tournefortii), Wireweed (Hogweed)	QLD, NSW ACT, Vic, TAS, WA Only  SA Only	1.4 Litres per hectare	
<b>Barley, Commercial Rye, Triticale</b>  From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)	Volunteer Peas	SA Only	1 Litre per hectare Plus 150 ml per hectare Dicamba 200 g/L	Apply when most peas are 5-7 cm high.
<b>Oats, Wheat</b> From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)			1 Litre per hectare Plus 300 ml per hectare Dicamba 200 g/L	
<b>Grass Pastures</b>	Bathurst Burr, Caltrop (Yellow Vine), Chicory, Crofton Weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora Burr, Pennygrass, Potato Weed, Ragwort, Ripplewort, Shepherd's Purse, Smartweed, Soft Rush, Stinkwort, Thistles including Cockspur, Scot (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All States	2.8-4 Litres per hectare  190-270 ml per 100 Litres  60 ml per 15 Litre Knapsack	Use the higher rate on larger weeds
	Cotton Thistle	TAS only		



a division of  
Amgrow Australia Pty Ltd

# Double Time - Post Emergent Selective Herbicide

**Application: (Refer to product label for more detailed instructions)**

CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
<b>Grass Pastures</b>	Buttercups, Hoary Cress Seedlings, Horehound, Khaki Weed, Purple-Top, Veined Verbena	All States	4-7 Litres per hectare 470 ml per 100 Litres 100 ml per 15 Litre Knapsack	When necessary a second treatment may be made after 4 weeks
<b>Turf</b>	Bindy-eye, Capeweed, Catsear, chickweed, Clovers, Cotula, Creeping Oxalis, Cudweeds, Dandelion, Dock, Fleabanes, Fumitory, Jo-Jo (Onehunga), Lamb's Tongue, Medic, Mustards, Narrowleaved Plantain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet Pimpernel, Slender Thistle, Soldier Thistle, Spear Thistle, Toad Rush, Variegated Thistle, Wild Turnip, Wireweed (Hogweed)	All States	<b>Golf Tee, Golf Green:</b> 100 ml in 15 litres water per 150m <sup>2</sup>  <b>Bowling Green:</b> 900 ml in 200 litres water per 1400m <sup>2</sup>  <b>Large Turf Area:</b> (Field, Park, Golf Fairway) 6.5 L in 250-400 L water per 10,000 m <sup>2</sup> (1 hectare)	Apply to actively growing weeds and moist soil.  DO NOT use on Lippia, Strawberry Clover or Buffalo Grass.  DO NOT mow turf for two days before or after application or fertilise within two weeks of spraying.
	Fat-Hen, Purple-Top, Red flowered Mallow, Staggerweed, Swinecress	QLD, NSW ACT, TAS SA, WA Only		
<b>Non-crop situations</b>	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All States	5.2 L per hectare 350 ml per 100 Litres 80 ml per 15 litre knapsack	A repeat spray may be necessary

