FarmAXE RLP Oct 2013

POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FarmAXE® 125 SC Insecticide

ACTIVE CONSTITUENT: 125 g/L BETACYFLUTHRIN

GROUP 3A INSECTICIDE

For the control of cockroaches, spiders, ants, fleas, flies, silverfish, bed bugs, mosquitoes, clothes moths, carpet beetles and stored product pests in the situations specified in the DIRECTIONS FOR USE

[IMPORTANT: Read the attached booklet before use] [IMPORTANT: Read this booklet before use]

Contents: 100 mL

GENERAL INSTRUCTIONS

Mixing

Prior to pouring, shake container vigorously, then add the required quantity of FarmAXE 125 SC Insecticide to water while stirring. **Application**

Do not spray the moving parts of machinery, electric motors or switch gear with this water-based product.

Note on application to carpets

Whilst carpets are regarded as porous surfaces, excessive liquid application may produce water-stains and therefore the non-porous surface rate should be adopted if possible. Hence for short pile carpet, the 5 L dilution rate (per 100 m²) should be sufficient. However, for long pile or shag carpet, the 10 L dilution rate (per 100 m²) may be required to ensure effective penetration and application to all of the carpet pile.

Since carpets vary in thickness, colour and material, it is recommended that a small, inconspicuous area of carpet be sprayed and allowed to dry first to ensure that staining will not occur. Allow treated carpets to dry before resuming activity on them.

PRECAUTIONS

For surface treatment only – do not spray into the atmosphere using foggers or other devices designed to create an aerosol. Use in food processing areas is limited to crack, crevice and barrier strip treatment only.

When using in animal houses, do not spray directly onto livestock, their food, water, litter, feed bins or surfaces that are accessible to the stock.

Do not spray directly on humans, exposed food, food utensils or food preparation areas.

Avoid contact of spray with timber to which a stain is subsequently to be applied, unless the owner is notified that the timber will need additional preparation, consistent with that if water alone had contacted the timber. (Evidence of water contact on timber is sometimes only seen once the stain has been applied).

BEFORE USE

Remove or cover all exposed foodstuffs. Cover all dishes and utensils and surfaces where food is prepared.

AFTER USE

Thoroughly ventilate treated areas. Clean up thoroughly before processing or serving resumes.

USE AS A RESIDUAL SURFACE TREATMENT FOR CONTROL OF STORED PRODUCT PESTS

The following measures should be observed:

Ensure food items in store are not exposed to spray. If necessary, remove food items from the area to be treated. Allow all treated surfaces to dry before replacing food items. Where possible avoid spraying actual surfaces which food is likely to directly contact. Do not allow spray to contact food packaging material.

Discard any foodstuff accidentally contaminated with spray solution.

Insecticide Resistance Warning

GROUP 3A INSECTICIDE

For insecticide resistance management FarmAXE is a Group **3A** insecticide. Some naturally occurring insect biotypes resistant to FarmAXE and other Group **3A** insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if FarmAXE or other Group **3A** insecticides are used repeatedly. The effectiveness of FarmAXE on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Sundew Solutions Pty Ltd accepts no liability for any losses that may result from failure of FarmAXE to control resistant insects. FarmAXE may be subject to specific resistance management strategies. For further information contact your local supplier, or Barmac representative.

FarmAXE RLP Oct 2013

DIRECTIONS FOR USE

SITUATION	RATE	PEST	CRITICAL COMMENTS
Meatworks			On non-porous surfaces, apply at a rate of 5 L of prepared spray per 100
(non-product			m ² not exceeding the point of run-off.
areas), farm			On porous surfaces or for use through power equipment, apply at a rate
buildings, food			of 10 L of prepared spray per 100 m ² not exceeding the point of run-off.
processing	Initial or	Cockroaches	Conduct a thorough inspection and apply spray to known and potential
establishments,	cleanout rate		cockroach harbourages. A barrier spray around areas where cockroach
shops, offices,	4 mL/L		infestations may be originating is also suggested. Repeat application as
factories,	Maintenance		required. Where application of a spray is not suitable, it is recommended that a registered cockroach gel be used.
storerooms.	rate	Web spinning	Spray spiders, their webbing and areas where spiders may hide. Leave
hospitals,	2 mL/L	spiders	webbing in place for a few days to optimise efficacy.
schools,	2 1112/2	Non-web	Spray areas where spiders are known to occur.
barracks,		spinning spiders	opray areas micro opiasio are known to coour.
houses, ships,		Ants	Where possible, locate the nest and spray directly into nest entrances.
rubbish tips			Penetration of insecticide can be assisted by pre-wetting sandy and
מקוז וופוטטוזו			porous soils with water. A thorough soaking treatment with insecticide
			mixture is required to destroy the nest. If nest cannot be located then
			spray infested areas.
		Fleas	Clean or vacuum infested areas and dispose of the collected dust in the
			garbage in a sealed plastic bag. Lightly spray all infested areas
			concentrating under furniture, etc. Treat pets with a registered flea
		Flice and	control product. See note on application to carpets.
		Flies and	Spray surfaces where flies and mosquitoes are known to come to rest. Flies – include external surfaces that are exposed to the morning sun.
		mosquitoes	Mosquitoes – spray walls, awnings, fences and vegetated areas near
			buildings.
			Note: the perception of the level of control of flying insects achieved
			within any given area treated with a residual surface spray will be affected
			by the extent of migration of individuals from external untreated areas. A
			residual spray will only control those insects that alight on the treated
			surfaces and therefore complete elimination of flying insects from the
			treated area should not necessarily always be expected.
		Silverfish	Spray areas of potential and existing infestations eg under skirting boards
			and furniture where dark conditions may favour silverfish populations. If
			spraying inside cupboards or storage areas, remove clothing and
		Clothes moths	utensils, etc before spraying.
		Ciotnes moths	Spray surfaces in areas where clothes are stored. DO NOT spray the clothes.
		Carpet beetles	Apply to infested carpets, particularly in areas of little or no traffic and
		Carpot bootios	underneath windows where adult beetles may be entering the building.
			See note on application to carpets.
		Bed bugs	Apply to the bed frame and walls, cracks, crevices and skirting boards in
			the vicinity of the bed.
		Stored product	The use of this product for the control of stored product pests should be
		pests	considered part of an integrated strategy to prevent development of
			insect populations. Best results will be obtained where good hygiene is
			maintained. Infested areas should be cleaned prior to application and
			regularly cleaned thereafter to reduce food sources for pests. When
			treating for beetle pests, pay special attention to sheltered areas such as
			cracks and crevices, under furniture and equipment, floors, walls and shelves. Adult moths can be controlled by spraying walls and surfaces
			where the flying moths come to rest.
Rubbish tips	Knockdown	Flies	Spray surfaces of rubbish. Apply 200-300 mL of spray/m².
	spray		The state of the s
	1 mL/2 L		
			ANY MANNED CONTRADY TO THIS LABEL LINESES

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION USE IN FOOD PROCESSING AREAS IS LIMITED TO CRACK, CREVICE AND BARRIER STRIP TREATMENT ONLY

FarmAXE RLP Oct 2013 3

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. Cover fish tanks before using. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Avoid contact with skin. Do not inhale vapour and spray mist. When preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half face-piece respirator. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, and rubber gloves. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, respirator and if rubber wash with detergent and warm water, and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. Remove from contaminated area. Apply artificial respiration if not breathing.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

LIMITATION OF LIABILITY

This product is designed only for the particular purposes indicated on this label. Failure to use this product strictly as directed may be illegal; prove dangerous; render the product ineffective and void any Warranty or Liability of Sundew Solutions Pty Ltd. So far as it is lawfully able to do so, Sundew Solutions Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

FarmAXE® is a Registered Trademark of Sundew Solutions Pty Ltd

Distributed by: BARMAC (A Division of Amgrow Pty Ltd) 82 Christensen Road Stapylton QLD 4207

Ph: 07 3802 5050 Email: sales@barmac.com.au www.barmac.com.au

APVMA Approval Number 66704/57710

DOM: BN: