

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 **Monopoly**[®]
Herbicide

ACTIVE CONSTITUENT: 720 g/L MSMA

GROUP Z HERBICIDE

**For the control of certain weeds in Cotton,
Sugarcane, Turf and non crop areas as
per the Directions For Use.**

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING




Amgrow

Distributed by: Amgrow Pty Ltd ABN 81 100 684 786

Unit B2A, 3-29 Birnie Avenue, Lidcombe NSW 2141 • TELEPHONE: 1800 631 008

**Monopoly Herbicide
DIRECTIONS FOR USE**

**Restrains: DO NOT cut, mow or cultivate treated areas that have been sprayed for at least 14 days as this may reduce the herbicide action.
DO NOT spray if rain is expected within 6 hours.**

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Cotton	Bathurst Burr Columbus Grass Johnson Grass Noogoora Burr Nut Grass Paspalum	Qld, NSW, WA only	GROUND APPLICATION BROADCAST: 3.1 litres/ha in 330 litres of water. BAND: 0.94 litres in 125 litres of water per sprayed hectare. (For 30 cm wide rows).	Spray young actively growing weeds under dry, hot conditions when air temperatures are above 21°C. DO NOT apply more than twice in one season. Where possible direct spray to base of cotton plant.
	Barnyard Grass Pig Weed Spiny Burr Grass Summer Grass Yellow Vine	Qld, NSW only	AERIAL SPRAYING: 3.1 litres/ha in 25-80 litres of water.	For non directed and aerial sprays a red discolouration and slight burning may occur. DO NOT apply on windy days. Commence spraying when cotton is 7-8 cm high and before first bloom. DO NOT spray after first bloom. Repeat 3 weeks later if necessary.
Sugar cane	Barnyard Grass Crab Grass/Summergrass Green Summergrass Itch Grass	Qld only	SPOT SPRAYING: 1.1 litres per 100 litres of water. Add a non ionic wetting agent.	Apply as a directed spray avoiding contact with sugar cane foliage as far as possible. Spray when sugar cane is 50-80 cm high. Add a non ionic wetting agent to improve the activity of this product. Whenever sugar cane is sprayed directly with product some visual damage may occur. Some sugar cane varieties are sensitive to herbicide spray damage and full cover spraying of these varieties may result in unacceptable crop phytotoxicity. Local varietal responses should be confirmed before using this product.
	Vasey Grass or Giant Paspalum	Qld, NSW only	6.6 litres/Ha in a minimum of 300 litres of water. SPOT SPRAYING: 1.1 litres in 100 litres of water. Add a non ionic wetting agent.	Apply as a directed spray minimising contact with sugar cane foliage. Spray when cane is 50-80 cm high. Add a non ionic wetting agent to improve the activity of this product. Best results are achieved if weeds are at the seedling stage and stools are small. Use sufficient spray volume to thoroughly wet weeds. Use higher volumes of water when weeds are large. Whenever sugar cane is sprayed directly with product some visual damage may occur. Some sugar cane varieties are sensitive to herbicide spray damage and full cover spraying of these varieties may result in unacceptable crop phytotoxicity. Local varietal responses should be confirmed before using this product.

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Non crop areas roadsides, fencelines & storage areas	Bathurst Burr Noogoora Burr Paspalum Johnson Grass Nut Grass Spiny Burr Grass Most Summer grasses	Qld, NSW, Vic, SA, WA only	11-13.3 litres/Ha in 500 litres of water for boom application or 1.1 litres per 100 litres of water for spot spraying.	Use lower rate for seedling weeds. Repeat 3-4 weeks later on established perennial weeds.
Control of Prickly Pear	Common Prickly Pear Devils Rope Drooping Prickly Pear Indian Fig Wheel Cactus	SA only	1.1 litres in 40 litres of water	Apply during Summer using a power sprayer when air temperature is above 30°C. Thoroughly cover all pad surfaces. Re-spray any re-growth that may occur.
Turf couch & hybrids of couch only (<i>Cynodon</i> spp.)	Crab Grass/Summergrass Paspalum	Qld, NSW only	6.6 litres/ha in 300 litres of water. Split applications of 3.3 litres/ha in 300 litres of water, applied 7-10 days apart will reduce phytotoxicity.	DO NOT apply to Bent, Qld Blue Couch (<i>Digitaria</i> spp.), Fescue, Blue Grass, Kikuyu or Dichandra lawns or turf as severe damage or death to the turf can occur. While some yellowing is expected this can be kept to a minimum by splitting the application to two applications 7-10 days apart. Weeds should be sprayed when young and actively growing. A repeat application in 14 days may be necessary. Avoid using at high temperatures as this may increase phytotoxicity. DO NOT mow for 4 days before or after application and do not water for at least 2 days after application.

NOT TO BE USED FOR ANY PURPOSE, OR IN MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS TO BE RECOMMENDED FOR USE IN THE HOME GARDEN.

WITHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCKFEED FOR 5 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

The use of a surfactant with this product is generally not required except where specified in the directions for use. For best results apply under hot, dry conditions when air temperatures exceed 25°C. As spray is absorbed mainly through the foliage thorough spray coverage of weeds is essential. Weeds should begin to yellow within 10 days of application.

MIXING

This product readily mixes with water. Half fill spray tank then add product with agitation. Fill tank.

CLEAN UP

Clean equipment by thoroughly washing with water.


PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish. DO NOT contaminate dams, rivers, drains or streams with chemical or used container.

RESISTANT WEEDS WARNING

Monopoly Herbicide is a member of the  Organoarsenicals group of herbicides. The product has the Herbicides with unknown and probably diverse sites of action mode of action. For weed resistance management, the product is a Group Z herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group Z herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group Z herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Amgrow Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

STORAGE AND DISPOSAL

Store in a locked room or place away from children, animals, food, feedstuff, seeds and fertilizers.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemical on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. May irritate the eyes and skin. Avoid contact with the eyes and skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which can be obtained from the supplier.

CONDITION OF SALE: Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions.

® Monopoly is a registered trademark of Amgrow Pty Ltd

APVMA Approval No.: 66365/53509



Distributed by:

Amgrow Pty Ltd (Commercial Products Division)
ABN 81 100 684 786
Unit B2A, 3-29 Birnie Avenue, Lidcombe NSW 2141
Telephone: 1800 631 008
www.amgrow.com.au