



a division of
Amgrow Australia Pty Ltd

Product Information Sheet

PCO Permethrin D

Analysis: 10 g/kg permethrin 25:75

Group **3A** Insecticide

Permethrin D is a professional insecticidal dust formulation for the control of cockroaches, ants, fleas, termites, european wasps and other insects in certain situations.

Benefits:

- An easy to handle and apply dusting formulation of the popular insecticide for controlling common insect pests in domestic, commercial and industrial situations
- Ideal for crack and crevice treatment of insect harborages
- Can be used to remove fleas from carpets indoors
- Reduced odour and photodegradation of active compared to liquid alternatives
- Can be applied by hand puffing applicator for specific purposes, refer to label



Pack Sizes: 10kg

Directions for application (refer to product label for more detailed instructions)

Situation	Pest	Rate	Comments
Domestic, industrial and institutional premises	Cockroaches, silverfish	10g/m ² applied with a powder blower	Blow powder into cracks and crevices. Attention should be given to areas where insects congregate ie. warm, dark areas and around heat sources (stoves, radiators, fireplaces etc).
	Ants		Sprinkle powder around nests and along routes. When ants are invading houses treat the areas where they are entering the premises.
	Fleas		Dust powder directly into carpets and brush into pile. Also treat areas where flea larvae may develop.
	Bedbugs		Apply to bedroom furniture and mattresses as well as to any cracks, crevices or areas where insects may hide
Roof voids, wall and floor voids, crawl spaces	Cockroaches, silverfish, ants, fleas, spiders and bird mites		Blow dust into all void areas, ensuring all surfaces are thoroughly treated. Repeat application if re-infestation occurs. Remove all birds nests and introduce physical bird barriers to prevent reinfestation of bird mites.
Mushroom beds	Mushroom sciarid flies		Dust mushroom beds when necessary. Do not apply more than twice per week. Discontinue use when mushrooms reach the size of a ten cent piece. Do not apply later than 3 days before harvest.
Skin and hide preparation	Hide beetles		Dust skins while on drying racks. Dust the edges of dried baled skins.
Seed potatoes	Potato moth	600g/T or 30g/50kg bag of seed potato	Apply dust while filling storage containers. Treated seed potatoes must only be used for sowing. Do not use as food or feed. Store treated seed potatoes away from other potatoes, food or feed. Mark bags or containers where treated seed potatoes are stored indication that the potatoes have been treated.

This publication is a guide only and no substitute for professional or expert advice. The product label should be consulted before use of any of the products referred to in this publication. Barmac (a division of Amgrow Australia Pty Ltd) shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein. Before using, always read the product label. More information: Contact your local Barmac distributor or go to www.barmac.com.au