



a division of
Amgrow Australia Pty Ltd

PRODUCT INFORMATION SHEET

Wettable Sulphur

Active Constituents: 800g/kg sulphur

Wettable sulphur is a preventative fungicide and miticide (Group Y) and is for the control of certain fungal diseases in grape vines, pome and stone fruit and vegetables. It is also for the control of mites in citrus, grape vines and vegetables

Benefits:

- Economical crop protection from powdery mildew, rust, brown rot, black spot and mites in registered crops
- Permit for use in dubosia (PER13863) for red spider mite
- A dry flowable formulation that disperses rapidly
- Can be applied in tank mixes with a wide range of other chemicals and/or fertilisers



Product Characteristics:

Colour	Specific Gravity
Brown/Yellow powder	1.8-2 g/cm ³

Pack Size:

Available in 2kg packs



Brown rot in stone fruit



Citrus bed mite damage in lemon



a division of
Amgrow Australia Pty Ltd

Wettable Sulphur

Directions for Application (Refer to product label for more detailed instructions)

CROP	WEEDS	STATE	RATE/100L WATER	CRITICAL COMMENTS
Vegetables (not cucumbers or rockmelons)	Powdery Mildew, Rust, Tomato Russet Mite, Bean Spider Mite, Two spotted Mite	SA, WA, VIC, TAS only	200 to 300g	Apply when pest or disease is first seen and repeat as necessary. Use the higher rate in crops with a large leaf cover or when conditions favour disease or mite development.
	Powdery Mildew Rust	QLD only	350g	
	Powdery Mildew, Rust, Tomato Russet Mite, Bean Spider Mite	NSW only	200g	
Citrus	Citrus Rust Mite, Brown Citrus Rust Mite, Citrus Bud Mite	NSW, WA, QLD, VIC & SA only	400 to 500g	Spray July to August with thorough coverage. Do not use on citrus in hot weather or within 21 days of an oil spray. Use the higher rate for heavy infestations.
	Citrus Bud Mite	TAS only	200 to 300g	
	White Louse Scale	NSW only	200 to 300g	May be added to copper spray at petal fall or applied separately during spring or autumn. Do not use on citrus in hot weather or within 21 days of an oil spray. Use the higher rate for heavy infestations.
Grapes	Grape leaf Blister Mite	NSW, VIC, TAS, SA & WA only	200 to 300g	Apply at 2-4 weekly intervals depending on the incidence of pest or disease. Use the higher rate for heavy incidence of infestation.
	Vine Mite	QLD only	350g	
	Powdery Mildew	NSW, VIC, SA, WA & TAS	200 to 300g	Apply when shoots are 20-25 cm long and repeat at intervals of 10 -14 days during humid weather. Use the higher rate early in the season before temperatures rise.
	QLD only	350g		
Stone Fruit (except apricots)	Rust	NSW, VIC, SA, WA, & TAS only	200 to 300g	Apply about 4 weeks after petal fall. Use the higher rate when conditions favour disease development.
		QLD only	350g	Apply at petal fall and then at intervals of 3-4 weeks until mid-late January.
	Brown Rot	NSW, VIC, TAS & SA	200g	Apply 4 weeks after petal fall then as cover sprays.
QLD only		350g	Apply at petal fall only.	
Pome Fruit	Powdery Mildew	NSW, VIC, TAS, SA & WA only	200 to 300g	Apply at pink bud, petal fall and then at intervals of 2-3 weeks. Use the lower rate if weather is hot and sensitive crops. Applying additional products in the same tank mix increases the phytotoxic risk to crops. Because climatic and soil conditions, application methods, irrigation and agricultural practices are beyond the control of Barmac and cannot be foreseen, Barmac accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.
		QLD only	350g	
	Black Spot (Scab)	VIC, TAS & SA only	200g	
		QLD only	350g	
Ornamentals	Powdery Mildew, Two Spotted Mite	VIC, TAS, SA, WA	200g	
		QLD only	350g	
	Powdery Mildew	NSW only	200g	

NOTE: Do not foliar apply during the heat of the day (> 25 DegC) when evaporation rates are at their highest. It is recommended that when foliar applying to a crop or area for the first time in combination with other chemicals, a small test area should be sprayed and observed for phytotoxicity prior to the total spray. Foliar spraying is recommended during early morning or late afternoon. Use the minimum foliar application rate on young or sensitive crops. Applying additional products in the same tank mix increases the phytotoxic risk to crops. Because climatic and soil conditions, application methods, irrigation and agricultural practices are beyond the control of Barmac and cannot be foreseen, Barmac accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.