

Thiram DG - Preventative Fungicide (Group Y) Active Constituent: 800 g/kg Thiram

For the control of certain fungal diseases of strawberries. Thiram is a member of the dithiocarbamate class of fungicides.

This group of fungicides has proven to be one of the most important and frequent control measures utilised today, they are primarily used as foliar protectants.

Due to their versatility, dithiocarbamates are able to provide excellent protection against a wide variety of fungal diseases with a low risk of resistance. The low resistance risk is related to thiram targeting multiple sites of action at a cellular level of the target disease complex.

Thiram DG Benefits

- Economical crop protection from a wide range of fungal diseases
- Thiram due to its multiple sites of action can be used as an effective resistance management tool when rotated with other mode of action fungicides
- A dry flowable micro granular formulation that dissolves rapidly
- Can be applied in tank mixes with a wide range of other chemicals and/or fertilisers



Pack Sizes: Available in 2kg and 20kg bucket

Product Characteristics

| Colour | Granule Size |
|----------------------|-----------------------|
| Brown Micro Granules | 0.6 g/cm ³ |



Black Spot in Strawberry



Grey mould in Strawberry

Directions for application (refer to product label for more detailed instructions)

| State | Disease | Rate | Critical Comments |
|----------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Queensland, Victoria, South Australia, Tasmania and Western Australia Only | Black spot (Colletotrichum acutatum), Grey Mould (Botrytis cinerea) | 150g/100 litres of water | Apply as a protective spray at flowering and then at 10-14 day intervals as necessary until 2 days before harvest. DO NOT exceed a maximum water volume of 1200 L/ha |
| COMPATIBILITY | Do not mix with lime sulphur or bordeaux sprays. DO NOT mix with fungicides containing copper. | | |

This publication is a guide only and no substitute for professional or expert advice. The product label should be consulted before use of any of the products referred to in this publication. Amgrow Specialty Ag shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein. Before using, always read the product label.