



# Key-El Iron

13% Iron as EDTA Chelate



**25kg** Net

# CAUTION: KEEP OUT OF REACH OF CHILDREN

## Key-El Iron

### GENERAL RECOMMENDATIONS

Key-El Iron is specifically formulated for enhanced delivery of chelated trace elements via soil, foliar application and fertigation.

Key-El Iron may be effectively applied with many agricultural products such as fungicides, insecticides, herbicides and other fertilisers. For unfamiliar tank mix combinations, sufficient evaluation to determine efficacy and crop safety may be warranted. Optimum rate of application will vary depending on crop variety, crop load and general crop health.

### GENERAL APPLICATION GUIDELINES:

- Do not apply a second foliar application within 14 days of the last application.
- Do not apply during flowering or fruit colour change.
- When applied with other Key-El products, the combined rate should not add up to more than maximum rate for each individual product.
- Do not mix with acid based products.

**MIXING INSTRUCTIONS:** Put 1/3 to 2/3 of total desired water volume in tank. Add desired amount of Key-El Iron and agitate until thoroughly mixed. Fill tank with remainder of desired water. A jar test is a good field practice for evaluating compatibility of multiple nutrient mixtures.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Store in a cool dry, locked area out of reach of children. Protect from excessive heat or cold. Dispose of container in accordance with local, state and federal regulations.

**CAUTION:** Do not swallow. Avoid contact with skin, eyes and clothing. May cause eye irritation. Wash hands thoroughly after using. In case of eye contact, flush eyes with water for at least 10 minutes and seek medical attention.

**NOTICE:** This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective.

Crops	Foliar Application per Ha	Foliar Water Rate	Soil Application per Ha
Cereals, Grains, Legumes, Oilseed, Cotton, Fodder	0.25 - 1.0 kg	1% or 100g per 10L of water	1 - 3 kg
Potatoes & Vegetables	0.5 - 1.5 kg	0.1% or 100g per 100L of water	4 - 7 kg
Grapevines (Wine and Table)	0.5 - 1.0 kg		4 - 8 kg
Fruit & Nut Tree Crops	0.5 - 1.0 kg		5 - 10 kg
Strawberries & other Berry Crops	0.1 - 0.5 kg		5 - 10 kg



Manufactured in China for  
Barmac, a division of Australian Agribusiness  
(Holdings) Pty Ltd  
ABN 61 135 355 958  
Suite 201, 3 Rider Boulevard, Rhodes NSW 2138  
Ph: (07) 3802 5050  
[www.barmac.com.au](http://www.barmac.com.au)

Batch No:

DOM:



9 322321 004255