



a division of Amgrow

NUTRAFEED FLOWER PRO - Liquid Fertiliser

NUTRAFEED LIQUID FLOWER PRO NPKS 7-15-10-0

Additional Macro and Micro Nutrients (%w/v)

Ca %	Mg %	B %	Zn %	Fe %	Mn %	Cu %	Mo %
-	-	0.11	0.14	-	-	0.1	-



NUTRAFEED LIQUID FLOWER PRO

Flower Pro is a new high analysis liquid NPK + TE fertiliser suitable for foliar or soil application on a wide range of horticultural and broad acre crops.

FLOWER PRO BENEFITS

- Significantly improves flowering uniformity
- Promotes early season flowering reduces late flowering incidence
- High phosphorus and potassium analysis to trigger flowering
- Zinc improves flower fertility, seed, fruit and tuber size
- Copper stimulates floral bud initiation
- Boron improves pollen viability and reduces bud and flower drop
- Chloride, sulphate, nitrate free formulation



K Deficiency in Citrus



P Deficiency in Strawberry



Zn Deficiency in Tomato




B Deficiency in Grapes

PRODUCT CHARACTERISTICS

Colour	pH	SG
Blue	6.8	1.37

PACK SIZES

Available in 20, 200 and 1000 L packs.





a division of Amgrow

NUTRAFEED FLOWER PRO - *Liquid Fertiliser*

APPLICATION

Flower Pro can be used as a foliar or soil applied supplement in a regular nutrition program for applicable crops. As a foliar spray add a wetting adjuvant and stick to the dilutions in the table below with a maximum rate of 7L product/ha. Multiple applications may be needed throughout the season. The application rate may need to be varied with changes in plant size, canopy or crop load.

DIRECTIONS FOR USE

Crop	Foliar (L/100L water)	Fertigation (L/ha)	Comments
Berry fruits	0.5-1	8-10	Apply lower rate to young plants. Apply leading up to and during flowering at 10-14 day intervals
Citrus	-	10-15	Apply leading up to and during flowering at 10-14 day intervals. Fertigation only
Cucurbits	0.5-1	8-10	Apply lower rate to young plants. Apply leading up to and during flowering at 10-14 day intervals
Broadacre (cotton beans, chickpea)	1	-	Apply in the lead up to flowering
Pomefruit / Stonefruit	-	10-15	Apply leading up to and during flowering at 10-14 day intervals. Fertigation only
Potato	0.5-1	8-10	Apply lower rate to young plants. Apply leading up to and during flowering at 10-14 day intervals
Tomato / Capsicum	0.5-1	8-10	Apply lower rate to young plants. Apply leading up to and during flowering at 10-14 day intervals
Tree Nut Orchards	0.25	10-15	Apply leading up to and during flowering at 10-14 day intervals
Tropical Fruit (avocado, banana, mango, pines etc)	0.25	10-15	Apply leading up to and during flowering at 10-14 day intervals
Vegetables / Brassica	0.5-1	8-10	Apply lower rate to young plants. Apply leading up to and during flowering at 10-14 day intervals
Vines/Grapes	0.5-1	8-10	Apply lower rate to young plants. Apply leading up to and during flowering at 10-14 day intervals

NOTE:

The suggested rates of application are designed for typical use conditions and should be used as a guide only. Do not foliar apply during the heat of the day (> 25 DegC) when evaporation rates are at their highest. It is recommended that when foliar applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed for phytotoxicity prior to the total spray. Foliar spraying is recommended during early morning or late afternoon. Use the minimum foliar application rate on young or sensitive crops. Applying additional products in the same tank mix increases the phytotoxic risk to crops. Because climatic and soil conditions, application methods, irrigation and agricultural practices are beyond the control of Barmac and cannot be foreseen, Barmac accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.