



a division of Amgrow

NUTRAFEED NITRO TRACE - *Liquid Fertiliser*

NUTRAFEED LIQUID NITRO TRACE NPKS 32-0-0-0.6

Additional Macro and Micro Nutrients (%w/v)

Ca %	Mg %	B %	Zn %	Fe %	Mn %	Cu %	Mo %
-	0.38	-	0.11	0.19	0.06	-	0.05



NUTRAFEED LIQUID NITRO TRACE

Nitro Trace is a new high analysis liquid nitrogen with trace element fertiliser suitable for foliar or soil application on a wide range of horticultural and broad acre crops.

NITRO TRACE BENEFITS

- Effectively provides a one shot growth boost to promote crop vigour
- Ideal for crop recovery from waterlogged situations by encouraging photosynthesis
- Molybdenum improves the nitrogen utilisation response
- Zinc for stem and leaf expansion support
- Magnesium, iron and manganese for chlorophyll production and increased photosynthesis



N Deficiency Tomato



N Deficiency in Wheat




Waterlogged Cotton Crop

PRODUCT CHARACTERISTICS

Colour	pH	SG
Green/Brown	4.2	1.22

PACK SIZES

Available in 20, 200 and 1000 L packs.





a division of Amgrow

NUTRAFEED NITRO TRACE - *Liquid Fertiliser*

APPLICATION

Nitro Trace can be used as a foliar or soil applied supplement in a regular nutrition program for applicable crops. As a foliar spray add a wetting adjuvant and stick to the dilutions in the table below with a maximum rate of 10L product/ha for horticultural crops. Multiple applications may be needed throughout the season. The application rate may need to be varied with changes in plant size, canopy or

DIRECTIONS FOR USE

Crop	Foliar (L/100L water)	Fertigation (L/ha)	Comments
Berry fruits	0.75-1	8-10	Apply when a general boost is required or to lift nitrogen levels
Broadacre (cereals, canola, corn, cotton, beans)	10-20	-	Apply during vegetative growth stage to lift nitrogen levels or for waterlogged crop recovery. Use lower rate on legumes
Citrus	0.25-0.5	15-20	Apply when a general boost is required or to lift nitrogen levels
Cucurbits	0.75-1	8-10	Apply during vegetative growth stage or to lift nitrogen levels
Pomefruit / Stonefruit	0.25-0.5	15-20	Apply post-harvest and when a general boost is required or to lift nitrogen levels
Potato	0.75-1	8-10	Apply during vegetative growth stage or to lift nitrogen levels
Tomato	0.75-1	8-10	Apply during vegetative growth stage or to lift nitrogen levels
Tree Nut Orchards	0.25-0.5	15-20	Apply post-harvest and when a general boost is required or to lift nitrogen levels
Tropical Fruit (avocado, mango, banana, pines etc)	0.25-0.5	15-20	Apply post-harvest and when a general boost is required or to lift nitrogen levels
Vegetables/Brassica	0.75-1	8-10	Apply during vegetative growth stage or to lift nitrogen levels
Vines / Grapes	0.25-0.5	8-10	Apply during vegetative growth stage or to lift nitrogen levels

NOTE:

The suggested rates of application are designed for typical use conditions and should be used as a guide only. Do not foliar apply during the heat of the day (> 25 DegC) when evaporation rates are at their highest. It is recommended that when foliar applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed for phytotoxicity prior to the total spray. Foliar spraying is recommended during early morning or late afternoon. Use the minimum foliar application rate on young or sensitive crops. Applying additional products in the same tank mix increases the phytotoxic risk to crops. Because climatic and soil conditions, application methods, irrigation and agricultural practices are beyond the control of Barmac and cannot be foreseen, Barmac accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.