

## **Nutrafeed 4**

NPKS: 19 - 2 - 17 - 8.3

Product Information Sheet

#### **Nutrafeed 4**

Nutrafeed 4 is a soluble crystalline blend formulated for fertigation (application through irrigation water) or foliar application. Fertigation provides maximum root uptake of the applied nutrients particularly when fed through trickle irrigation. Application through trickle enables the nutrients to be placed in the area of the most intensive root activity. Foliar fertilisation is not a substitute for soil fertilisation, but a supplement to it. Nutrafeed 4 is a completely soluble and chloride free formulation, providing a complete range of nutrients necessary for optimal plant growth. NutraFeed 4 is a well balanced NPK soluble fertiliser blend with good levels of nitrogen, phosphorus, potassium, magnesium and trace elements for general growth and is suitable for a wide range of horticultural or broad acre crops at any growth stage

### **Nutrafeed 4 Benefits**

- Significantly improves overall crop fertility levels and increases general crop vigour
- Effectively provides a one shot multi nutrient boost to promote balanced growth
- Can be used at any crop stage due to its unique and balanced NPK ratio
- Contains Lo Biuret Urea, which is safer on plants than standard Urea
- Contain trace elements in chelated form maximizing plant uptake
- Chloride free formulation.

Solubility is 250g per 1L water

### **Additional Macro and Micro Nutrients**


Ca %	Mg%	В%	Zn %	Fe %	Mn %	Cu %	<b>Mo</b> %
-	0.3	0.01	0.01	0.08	0.04	0.01	0.002



**Pack Sizes** 

Available in 25kg poly bag.





N Deficiency in Tomato

K Deficiency in Banana



K Deficiency in Grape Vine



Amgrow Specialty Agriculture (A division of Australian Agribusiness (Holding) Pty Ltd



# **Nutrafeed 4**

NPKS: 19 - 2 - 17 - 8.3

Product Information Sheet

### **Application**

Nutrafeed 4 can be used as a foliar or soil applied supplement in a regular nutrition program for applicable crops. As a foliar spray add a wetting adjuvant such as Indicate 5 for best results. Multiple applications may be needed throughout the season. The application rate may need to be varied with changes in plant size, canopy or crop load. See your local agronomist for a prescribed nutrient budget to determine the optimum application rate for your specific purpose.

### **Directions for Use**

Crop	Foliar Kg/100L Water	Fertigation (kg/ha)	Comments	
Berry fruits	0.5-1	8-10	Apply when a general boost is required or regularly at 14 day intervals	
Broadacre (cotton, beans, chickpea)	0.5-1	-	Apply when a general boost is required	
Citrus	-	10-15	Apply when a general boost is required or regularly at 14 day intervals. Fertigation only	
Cucurbits	0.5-1	8-10	Apply when a general boost is required or regularly at 14 day intervals	
Pomefruit / Stonefruit	-	10-15	Apply when a general boost is required or regularly at 14 day intervals. Fertigation only	
Potato	0.5-1	8-10	Apply when a general boost is required or regularly at 14 day intervals	
Tomato	0.5-1	8-10	Apply when a general boost is required or regularly at 14 day intervals	
Tree Nut Orchards	0.25	10-15	Apply when a general boost is required or regularly at 14 day intervals	
Tropical Fruit (avocado, mango, banana, pines etc)	0.25	10-15	Apply when a general boost is required or regularly at 14 day intervals	
Vegetables / Brassica	0.5-1	8-10	Apply when a general boost is required or regularly at 14 day intervals	
Vines / Grapes	0.5-1	8-10	Apply when a general boost is required or regularly at 14 day intervals	

### **Note**

The suggested rates of application are designed for typical use conditions and should be used as a guide only. Do not foliar apply during the heat of the day (> 25 DegC) when evaporation rates are at their highest. It is recommended that when foliar applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed for phytotoxicity prior to the total spray. Foliar spraying is recommended during early morning or late afternoon. Use the minimum foliar application rate on young or sensitive crops. Applying additional products in the same tank mix increases the phytotoxic risk to crops. Because climatic and soil conditions, application methods, irrigation and agricultural practices are beyond the control of Amgrow Specialty Ag and cannot be foreseen, Amgrow Specialty Ag accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.

