



POISON

For Heavy Infestations

Single Feed

With Brodifacoum

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TOMCAT II RODENTICIDE



ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

Ready to use bait for use in commercial, agricultural and domestic buildings.



DIRECTIONS FOR USE

RESTRAINTS:

DO NOT exceed 3m between bait stations for mice or 9m for rats. DO NOT place bait stations in open. Mice require several bait stations, 50g per station (less than 1 g will kill a mouse). Rats require large quantities of bait in one or two locations, 50 - 100g per station (4 - 6g will kill a rat). For effective control continue until feeding ceases.

DO NOT apply bait to crops.

Baiting Strategy:

Eliminate as far as practicable, all other food sources. Bait infested area. Place bait stations under cover. Inspect bait stations daily. Replace eaten baits. If quickly eaten, increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

SITUATION: In and around agricultural, commercial and domestic buildings.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

Pellets are not subject to wind dispersal. Most deaths occur 4 to 7 days after consumption of bait. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Record the number and locations of bait stations for purpose of replenishment and later removal.

POISON

TOMCAT II RODENTICIDE



KILLS RATS & MICE

PRECAUTION: KEEP OUT OF REACH OF CHILDREN
Keep away from food and water and surfaces which come in contact with food and water.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, streams, rivers, waterways and drains with the bait or its used container. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. The product is restricted to use in and around buildings (within 2 metres), or enclosed spaces, e.g. drains. Search for and burn or bury all rodent bodies, do not place in refuse bins or rubbish tips. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.

Net 2.5 kg

TOMCAT is a trademark of Bell Laboratories, Inc., USA

STORAGE AND DISPOSAL

Store in a tightly closed, original container in a dry well ventilated area, as cool as possible out of a direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT inhale dust. DO NOT touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (phone Australia 131126; New Zealand 0800 764 766).

MSDS

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

The use of TOMCAT II is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

060607

APVMA APPROVAL NO: 51280/2.5KG/0407

Batch No.:

D.O.M.:

Manufactured by:
Bell Laboratories, Inc.
Madison, Wisconsin 53704 U.S.A.
www.belllabs.com

Net 2.5 kg

Australian agent:
w e b c o t
Level 1, APP House, 14 Rodborough Road
Frenchs Forest NSW, 2086, Australia
Phone: (02) 9984-2255 Fax: (02) 9984-2222
EMERGENCY CONTACT NO. (02) 9984-2255

51280_41694_2.5KG_LABEL_MPL_V2

Total area on A3 dieline:
52.3875cm x 12.7cm

75% to actual 100% size

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

For Heavy
Infestations

Single Feed

With
Brodifacoum



TOMCAT II

RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM
Contains Denatonium Benzoate (Bittering Agent)

Ready to use bait for use in commercial, agricultural and domestic buildings.

KILLS RATS & MICE

Manufactured by:
Bell Laboratories, Inc.
Madison, Wisconsin 53704 U.S.A.
www.belllabs.com

Net 10 kg

Australian agent:
w e b c o t
Level 1, APP House, 14 Rodborough Road
Frenchs Forest NSW, 2086, Australia
Phone: (02) 9984-2255 Fax: (02) 9984-2222
EMERGENCY CONTACT NO. (02) 9984-2255



RESTRAINTS:
DO NOT exceed 3m between bait stations for mice or 9m for rats.
DO NOT place bait stations in the open. Mice require several bait stations, 50g per station (less than 1g will kill a mouse). Rats require large quantities of bait in one or two locations, 50 - 100g per station (4 - 6g will kill a rat). For effective control continue until feeding ceases.
DO NOT apply bait to crops.

Baiting Strategy:
Eliminate as far as practicable, all other food sources. Bait infested area. Place bait stations under cover. Inspect bait stations daily. Replace eaten baits. If quickly eaten, increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

SITUATION: In and around agricultural, commercial and domestic buildings.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.


THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS
Pellets are not subject to wind dispersal. Most deaths occur 4 to 7 days after consumption of bait. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Record the number and locations of bait stations for purpose of replenishment and later removal.

POISON

TOMCAT II

RODENTICIDE



KILLS RATS & MICE

Net 10 kg

TOMCAT is a trademark of Bell Laboratories, Inc., USA

PRECAUTION: KEEP OUT OF REACH OF CHILDREN
Keep away from food and water and surfaces which come in contact with food and water.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
DO NOT contaminate dams, streams, rivers, waterways and drains with the bait or its used container. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. The product is restricted to use in and around buildings (within 2 metres), or enclosed spaces, e.g. drains. Search for and burn or bury all rodent bodies, do not place in refuse bins or rubbish tips. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.

SAFETY DIRECTIONS
Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre (phone Australia 131126; New Zealand 0800 764 766).

MSDS
For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE
The use of TOMCAT II is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA APPROVAL NO: 51280/10KG/0407 052507
Batch No.:
D.O.M.:

51280_41694_10KG_LABEL_MPL_V2

Total Area on A3 dieline:
80.3275cm x 19.3675cm

50% to actual 100% size