

# POISON

- **Controls Species Resistant To Warfarin**
- **Single Feed**
- **With Brodifacoum**



KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



*All-Weather*  
**BLOX**  
**Rodenticide**



ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM  
Contains Denatonium Benzoate (Bittering Agent)

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings.

## KILLS RATS & MICE

Manufactured by:

**Bell**  
LABORATORIES, INC.  
[www.belllabs.com](http://www.belllabs.com)  
Madison, WI 53704 U.S.A.  
MADE IN USA

CONTROLS SPECIES RESISTANT TO WARFARIN

PRIMARY PACK

CONTAINS 20 g RODENT BLOCKS

BLOCKS ARE ILLEGAL TO SELL SEPARATELY

DO NOT DESTROY OUTER CONTAINER WHILE  
PRODUCT STILL REMAINS

**Net 8 kg**

Australian Agent:

**Barmac**  
A Division of Amgrow Australia Pty Ltd  
(ABN 22 069 900 456)  
82 Christensen Road, Stapylton, QLD 4207  
EMERGENCY CONTACT NO. 1800 AMGROW  
(1800 264 789)  
[www.barmac.com.au](http://www.barmac.com.au)

APVMA APPROVAL NO: 50697/8KG/0207

**DIRECTIONS FOR USE****RESTRAINTS:**

DO NOT exceed 3m between bait stations for mice or 9m for rats.

DO NOT place bait stations in the open.

DO NOT apply bait to crops.

Situation	Pest	Rate	Critical Comments
In and around agricultural, commercial, domestic and industrial buildings	Rats ( <i>Rattus rattus</i> , <i>Rattus norvegicus</i> ) Mice ( <i>Mus musculus</i> )	1 to 3 blocks per bait station depending on level of infestation	Set out baits in places frequented by mice and rats but not more than 2m away from buildings etc. For Rats: Place 1 to 3 blocks at intervals of 5 to 9 m in infested areas. For Mice: Place single blocks at 2 to 3 m intervals in infested areas.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.**

**GENERAL INSTRUCTIONS**

Keep away from food and water, and surfaces that come in contact with food and water. Contains human taste deterrent, which prevents accidental consumption by children.

**Baiting Strategy:** Eliminate as far as practicable, all other food sources. Bait infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimized. Record the number and locations of bait stations for purpose of replenishment and later removal.

# POISON



## KILLS RATS & MICE

**GENERAL INSTRUCTIONS** (Continued from left panel)

Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased. The use of this product for the control of protected native rodents requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 meters) or in enclosed spaces, e.g. drains.

**Most deaths occur 4 to 7 days after consumption of bait.**

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**  
The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. Place baits in locations inaccessible to other animals or birds. DO NOT contaminate dams, streams, rivers, waterways and drains with the bait or its used containers. Look for dead rats or mice – burn or bury them, do not place in refuse bins or rubbish tips.

**Net 8 kg**

**MADE IN USA**

TOMCAT is a trademark of Bell Laboratories, Inc., USA

**STORAGE AND DISPOSAL****KEEP OUT OF REACH OF CHILDREN**

Store in a tightly closed, original container in a dry well-ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.

**SAFETY DIRECTIONS**

Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K<sub>1</sub> (Phytomenadione) is antidotal.

**FIRST AID**

If poisoning occurs, contact a doctor or the Poisons Information Centre (phone Australia 131126).

**SDS**

For further information please refer to the Safety Data Sheet which can be obtained from the supplier.

In a transport emergency dial 000, Police or Fire Brigade.

For specialist advice in an emergency only, call (02) 9037 2994 (24 h).

APVMA APPROVAL NO: 50697/8KG/0207

**GHS Hazard and Precautionary Statements**

**May cause damage to organs through prolonged or repeated exposure.**

Do not eat, drink or smoke when using this product. Avoid release to the environment. If eye irritation persists: seek medical attention. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. In case of fire: Preferred extinguishing media are carbon dioxide, dry chemical, foam, water fog.

013020

Batch No.:

D.O.M.:

Product Code: 933001

