

Totril

Selective Herbicide (Group C)
Analysis: 250g/L ioxynil

For the control of certain broadleaf weeds in onion crops.

Totril Benefits

- Trusted original Totril formulation
- Now available in a 10L container • Apply to onions between the three and eight leaf stage
- Target weeds from cotyledon to 6 leaf stage
- No withholding period if used as directed



Pack Sizes: Available in 10L packs.

Directions for application (refer to product label for more detailed instructions)

Situation	Weeds Controlled	Rate	Comments
Onions	Bellvine Burr Medic Capeweed Chickweed Climbing Buckwheat Common Heliotrope Common Sowthistle (Milk Thistle) Corn Gromwell (Sheepweed) Dandelion Dead Nettle (Henbit, Stingless Nettle) Fat-Hen Fumitories Green Amaranth Green Crumbweed (Keeled Goosefoot) Lesser Swinecress (Bittercress) Ox Tongue Pigweed (Portulaca) Potato Weed (Yellow Weed) Prickly Paddy Melon Saffron Thistle Scarlet Pimpernel or Blue Pimpernel Shepherd's Purse Slender Celery Smallflower Mallow (Marshmallow) Stagger Weed Threecornered Jack (Spiny Emex) Threeflower Nightshade Turnip Weed Wards Weed Wild Radish Wild Turnip Wireweed (Hogweed)	2.1 - 2.8L	<p>CROP: Apply between the three and eight leaf stage. Slight scorch or symptoms of wilting may occur but this is transient as subsequent growth and yield is unaffected.</p> <p>WEEDS: Apply between the cotyledon to 6 leaf stage. Most weeds will be more susceptible at the young plant stage than at later stages of growth. Use the higher rate when weeds are more mature.</p> <p>* Burr Medic, Capeweed, Saffron Thistle, Slender Celery, Smallflower Mallow and Wireweed will be controlled only in the cotyledon stage.</p>
		2.1L	Where Wireweed is dominant or weeds are more mature, 2 applications of 2.1 L/ha should be made at an interval of 14 to 21 days.

GENERAL INSTRUCTIONS

Mixing

To ensure even mixing, half fill the spray tank with clean water, add the required amount of product and stir thoroughly. Add the remainder of the water and mix again before spraying.

Application

The use of this product should follow standard pre-emergence herbicide treatments or cultivation, otherwise the weeds are likely to be too large for successful treatment by the time the onions have reached the tolerant stage. DO NOT apply on a windy day. DO NOT apply by aircraft or misting machine.

Boom Sprayer

Equip the machine with flat-fan nozzle tips designed to apply 440 or more litres of water per hectare at a pressure of 200 to 280 kPa. Calibrate the machine before use to ensure that the correct application rate is obtained. Check the height of the boom to ensure that the spray is evenly distributed.

Compatibility

As formulations of other manufacturers' products are beyond the control of Amgrow Specialty Ag (A division of Australian Agribusiness (Holdings) Pty Ltd), all mixtures should be tested prior to mixing commercial quantities.

