

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING



BARMAC

SureFlow Borax

Trace Element

ACTIVE CONSTITUENT:
110g/kg Boron as borax



For the treatment of
boron deficiency in fruit
and vegetables crops,
seedlings and hydroponics.
Can be foliar or soil
applied as outlined in the
directions of use.



Batch
Number:

Date of
Manufacture:

Net Contents: 1 kg

Amgrow Specialty Agriculture

(A division of Australian Agribusiness (Holding) Pty Ltd)

ABN: 61 135 355 958

Suite 201 Level 2,3 Rider Boulevard, Rhodes NSW 2138

www.amgrow-ag.com.au | 1800 063 619

SureFlow Borax

Trace Element

Directions of use:

<i>Crop</i>	<i>Application</i>	<i>Rates</i>	<i>Comments</i>
Fruit trees	Foliar	100-200g/100L	Apply in sufficient water to wet foliage. Apply in early crop life.
Vegetables		200-500g/100L	
Fruit trees	Soil	5-30kg/ha	Apply when deficiency symptoms are first noticed.
Vegetables		1.0-2.5kg/ha	Incorporate into soil prior to planting.
Turf	Foliar	3g/100m ²	Apply at commencement of growing season.
Ornamentals		400g/100L	Apply in sufficient water to wet foliage.

Note: Boron can be toxic when levels become too high. A leaf or soil test should be conducted to determine exact requirements necessary for optimum plant growth.

Notice:

This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Barmac will not accept any responsibility whatsoever and howsoever arising whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.