

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING



BARMAC

SureFlow Boric Acid

Trace Element

ACTIVE CONSTITUENT:
170g/kg Boron (B) as Boric Acid



For the treatment of boron deficiency in fruit and vegetables crops, citrus, stone fruit, pome fruit, turf and ornamentals.



Can be foliar or soil applied as outlined in the directions of use.



Batch
Number:

Net Contents: 1 kg

Amgrow Speciality Agriculture

a division of Australian Agribusiness (Holdings) Pty Ltd

ABN 61 135 355 958

Tel: (07) 3802 5050

PO Box 3056, Rhodes NSW 2138

Website: www.amgrow-ag.com.au

Date of
Manufacture:



9 322321 000356

SureFlow Boric Acid

Trace Element

Directions of use:

<i>Crop</i>	<i>Application</i>	<i>Rates</i>	<i>Comments</i>
Fruit trees	Foliar	100-200g/100L	Apply in sufficient water to wet foliage. Apply in early crop life.
Vegetable crops		200-500g/100L	
Fruit trees	Soil	5-30kg/ha	Apply when deficiency symptoms are first noticed.
Vegetables		1.0-2.5kg/ha	Incorporate into soil prior to planting.
Turf	Soil	3g/100m ²	Apply at commencement of season.
Ornamentals	Foliar	400g/100L	Apply in sufficient water to wet foliage

Note: Boron can be toxic when levels become too high. A leaf or soil test should be conducted to determine exact requirements necessary for optimum plant growth.

Notice:

This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Amgrow Specialty Agriculture will not accept any responsibility whatsoever and howsoever arising whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.