KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BARMAC

SureFlow Copper Sulphate Trace Element

ACTIVE CONSTITUENT:

250g/kg Copper as copper sulphate 125g/kg Sulphur as copper sulphate

For the treatment of copper deficiency in fruit and vegetables crops, citrus, stone fruit, pome fruit, turf and ornamentals.
Can be foliar or soil applied as outlined in the directions of use.









Batch Number:

Date of Manufacture: Net Contents: 1 kg

Amgrow Specialty Agriculture

a division of Australian Agribusiness (Holdings) Pty Ltd

ABN 61 135 355 958 PO Box 3056, Rhodes NSW 2138

Tel: (07) 3802 5050 Website: www.amgrow-ag.com.au

Directions of use:

Crop	Application	Rates	Comments
General rate	Foliar	200-500g/100L	Apply in sufficient water to wet foliage. Apply in early crop life.
Fruit trees	Soil	5-10kg/ha	Apply when deficiency symptoms are first noticed.
Vegetables		25-50kg/ha	Incorporate into soil prior to planting.
Turf	Soil	25-50g/100m ²	Apply at commencement of season.

Note: Copper can be toxic when levels become too high. A leaf or soil test should be conducted to determine exact requirements necessary for optimum plant growth.

Notice:

This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Amgrow Specialty Agriculture will not accept any responsibility whatsoever and howsoever arising whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.