KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BARMAC



SureFlow Moly

Trace Element

ACTIVE CONSTITUENT: 390g/kg Molybdenum as sodium molybdate

For the treatment of molybdenum deficiency in fruit and vegetables crops, seedlings and hydroponics. Can be foliar or soil applied as outlined in the directions of use.









Batch Number:

Date of Manufacture: Net Contents: 1 kg

Amgrow Specialty Agriculture

a division of Australian Agribusiness (Holdings) Pty Ltd

ABN 61 135 355 958 PO Box 3056, Rhodes NSW 2138

Tel: (07) 3802 5050 Website: www.amgrow-aq.com.au

SureFlow Moly

Trace Element

Directions of use:

Сгор	Application	Rates	Comments
General rate	Foliar	50-100g/100L	Apply in sufficient water to wet foliage. Apply in early crop life.
Seedlings	Foliar	40g/100L	Apply prior to transplanting
Hydroponics	2.5ppm can be achieved by mixing 7g of sodium molybdate per 1000L of water		

Note: Molybdenum can be toxic when levels become too high. One spray per crop is generally sufficient, except where deficiency is noted. Susceptible crops such as brassicas and cucurbits may require two sprays three weeks apart.

PROTECTION OF LIVESTOCK: Excessive molybdenum can be harmful to stock - do not apply fertiliser containing molybdenum on more than 25% of farm annually. Do not graze for four weeks after application.

Notice:

This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Amgrow Specialty Agriculture will not accept any responsibility whatsoever and howsoever arising whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.